



UNIVERSITY OF THE PHILIPPINES
Master of Science in Environmental Science

EDWARD P. CAJUCOM

**LOCALIZED SPATIOTEMPORAL DATA ANALYSIS ON THE EFFECT OF LAND
COVER CHANGES ON TEMPERATURE TRENDS OVER THE PHILIPPINES**

Thesis Adviser:

GAY JANE P. PEREZ, Ph.D.
Institute of Environmental Science and Meteorology
University of the Philippines Diliman

Thesis Co-adviser:

MARCELINO Q. VILLAFUERTE II, Ph.D.
Climatology and Agrometeorology Division
Philippine Atmospheric, Geophysical and Astronomical Services Administration

Date of Submission:

April 2021

Thesis Classification:

P

(This thesis is not available to the public. Please ask the library for assistance.)

ABSTRACT

LOCALIZED SPATIOTEMPORAL DATA ANALYSIS ON THE EFFECT OF LAND COVER CHANGES ON TEMPERATURE TRENDS OVER THE PHILIPPINES

EDWARD P. CAJUCOM

University of the Philippines, 2021

Adviser: **Dr. Gay Jane P. Perez**

Co-adviser: **Dr. Marcelino Q. Villafuerte II**

Reader: **Dr. Cherry L. Ringor**

The effects and impacts of global warming are now evident worldwide. These impacts are expected to exacerbate and this will highly affect the economy and therefore, the lives of the people. Also, as the population growth continues to expand, the growing needs of the population especially in terms of space resources are also increasing. However, this land conversion or removal of natural vegetation producing low-reflective, impervious, non-evaporating surfaces significantly affects the local weather and climate present in an area.

As observed from the ground, the Philippines has a mean temperature increase of 0.73°C during the 1951 – 2019 observation period. Remarkable changes in mean minimum temperature were observed as compared to mean maximum and mean temperature in several synoptic stations of the Philippine Atmospheric, Geophysical and Astronomical Services Administration (PAGASA) in the country. Also, historical analysis on the diurnal temperature variations exhibits high increase in minimum (0.013°C increase per year) compared to maximum (0.008°C) and mean (0.010°C) temperature.

As seen from the space, results of the satellite-based temperature trends indicate consistent findings with those observed by PAGASA. Country-wide statistical analyses on the historical observations and correlations of Land Surface Temperature (LST), Normalized Difference Vegetation Index (NDVI), and Normalized Difference Built-up Index (NDBI) show varying patterns of trends during the entire study period (1988-2019). Daytime historical LST trends show relative likelihood and significant correlations with PAGASA observed data, especially on mean temperature observations. Also, in terms of LST, the urban and highly populated provinces of Metro Manila ($+0.43^{\circ}\text{C}$), Pampanga ($+0.21^{\circ}\text{C}$), Nueva Ecija ($+0.20^{\circ}\text{C}$), Pangasinan ($+0.18^{\circ}\text{C}$), and Tarlac ($+0.17^{\circ}\text{C}$) all show high values of temperature change during the study period (1988-2019). Additionally, the analysis of the pixel-based vegetation trends using NDVI provides helpful contributions on the spatial extent and current vegetation cover of the country. However, albeit good correlation observed between LST & NDVI profiles, it was noted that it is still important to investigate the degree of land use structure complexity that is present and can be determined using NDBI. The spatial variation of land surface temperature is largely dependent on the spatial distribution of impervious surface coverage and spectral response of vegetation profile. Hence, this study was able to come up with a provincial level built-up map that is consistent with the findings observed on LST and vegetation cover. Similar to the observation noted in LST, the areas with highest mean NDBI throughout the study period are the provinces of Metro Manila (-0.04), Pangasinan (-0.14), Tarlac (-0.16), Pampanga (-0.19), and Nueva Ecija (-0.20).

Drawing upon the concept of disaster risk reduction and climate change adaptation, this work successfully developed an algorithm and utilized Google Earth Engine as a main tool to process and investigate the effect of urbanization in many cities and provinces of the country. The three case studies conducted in Metro Manila, Pampanga, and Romblon proved that NDBI is an accurate indicator of urbanization and can be used as a complementary metric to the traditionally applied NDVI especially in studies related to local land-use changes vis-à-vis urban development. However, the reliability of this method is lowered in mapping peripheral urban areas where barren or fallow land is widespread or in areas where exposure to extreme weather events is high (e.g. islands). Thus, several strategies and ways forward related to the algorithms developed by this study were recommended. Also, holistic lists of science-based plans and programs that are based on the findings of this study and on the current gaps and challenges in dealing with urban heat island and climate change adaptation planning were also given.